

MIP-1 alpha/CCL3, Human

Cat.no. PK0169

Product size: 10ug 50ug 1mg

Source: CHO

Species: Human

Biological Activity: ED50 < 100 ng/ml, measured in a calcium flux assay using CHO/Gα15 cells expressing CCR5.

Molecular Weight: 8-10 kDa, observed by reducing SDS-PAGE.

Formulation: Lyophilized after extensive dialysis against PBS.

Reconstitution: Reconstituted in ddH2O or PBS at 100 µg/ml.

Purity: > 95% as analyzed by SDS-PAGE and HPLC.

Endotoxin Level: < 0.2 EU/µg, determined by LAL method.

Storage: Lyophilized recombinant Human MIP-1 Alpha remains stable up to 6 months at lower than -70°C from date of receipt. Upon reconstitution, Human MIP-1 Alpha should be stable up to 1 week at 4°C or up to 2 months at -20°C.

Description: MIP-1 Alpha, also known as CCL3, G0S19-1 and SCYA3, is a small inducible monokine belonging to the intercrine beta (chemokine CC) family. It binds to CCR1, CCR4 and CCR5, and participates in the host response to invading pathogens by regulating the trafficking and activation of inflammatory cells, such as macrophages, lymphocytes, NK cells and dendritic cells. MIP-1 alpha polymorphisms are associated with HIV susceptibility or resistance. Recombinant MIP-1 alpha induces a dose-dependent inhibition of HIV and SIV infection.

Amino Acid Sequence:

00001 ADTPTACCFS YTSRQIPQNF IADYFETSSQ CSKPGVIFLT

00041 KRSRQVCADP SEEWVQKYVS DLELSA

Synonyms: C-C motif chemokine 3, G0/G1 switch regulatory protein 19-1, Macrophage inflammatory protein 1-alpha, MIP-1-alpha, PAT 464.1, SIS-beta, Small-inducible cytokine A3, Tonsillar lymphocyte LD78 alpha protein, MIP-1-alpha(4-69), LD78-alpha(4-69), CCL3, G0S19-1, MIP1A, SCYA3, LD78α, MIP1α

Note: For research use only, not for use in diagnostic procedure.